

# PHOTOSYNTHETIC HYDROGEN AND OXYGEN PRODUCTION BY GREEN ALGAE

E. Greenbaum and J. W. Lee

Oak Ridge National Laboratory  
P. O. Box 2008  
Oak Ridge, TN 37831

## Summary

Photosynthesis research at Oak Ridge National Laboratory is focused on hydrogen and oxygen production by green algae in the context of its potential as a renewable fuel and chemical feed stock. Beginning with its discovery by Gaffron and Rubin in 1942, motivated by curiosity-driven laboratory research, studies were initiated in the early 1970s that focused on photosynthetic hydrogen production from an applied perspective. From a scientific and technical point of view, current research is focused on optimizing net thermodynamic conversion efficiencies represented by the Gibbs Free Energy of molecular hydrogen. The key research questions of maximizing hydrogen and oxygen production by light-activated water splitting in green algae are (1) removing the oxygen sensitivity of algal hydrogenases; (2) linearizing the light saturation curves of photosynthesis throughout the entire range of terrestrial solar irradiance — including the role of bicarbonate and carbon dioxide in optimization of photosynthetic electron transport and (3) constructing real-world bioreactors, including the generation of hydrogen and oxygen against workable back pressures of the photoproduct gases.

## Introduction

Photosynthetic hydrogen production by green algae was discovered in the pioneering experiments of Gaffron and Rubin (1942). This work was followed up by Gaffron and his colleagues in a series of seminal papers (Gaffron, 1960; Kaltwasser *et al.*, 1969; Stuart and Gaffron, 1971 & 1972; ) as well as many others. From the point of view of renewable fuels and chemical feedstock production, it is light-activated simultaneous photoproduction of hydrogen and oxygen that is of primary interest. The pioneering experiments in this field were performed by Spruit (1958) who developed a novel two-electrode polarographic technique for the simultaneous measurement of hydrogen and oxygen transients by the green alga *Chlorella*. The principle conclusion he came to was that hydrogen and oxygen metabolisms are closely related and they derived from water splitting. Later work by Bishop and Gaffron (1963) indicated that light-dependent evolution of hydrogen appeared to require both photosystems.

Research on photosynthetic hydrogen production as a renewable energy source began in the 1970s (Gibbs, *et al.*, 1973; Lien and San Pietro, 1975; Mitsui *et al.*, 1977). Using the two-electrode technique Bishop *et al.* (1977) measured and interpreted hydrogen and oxygen production from a large group of green algae. However, due to the buildup of hydrogen and oxygen, with subsequent inhibition (*vide infra*) these reactions could be followed for only several minutes. Using a flow system that removed inhibitory oxygen, it was shown (Greenbaum, 1980) that sustained simultaneous photoproduction of hydrogen and oxygen could be observed for hours. In prior experiments using a glucose-glucose oxidase trap, Benemann *et al.* (1973) demonstrated hydrogen production from water by a chloroplast-ferredoxin-hydrogenase system. Measurement of the hydrogen analog of the Emerson and Arnold photosynthetic unit size (Greenbaum, 1977a, b) indicated that photogenerated reductant expressed as molecular hydrogen was derived from the mainstream of the photosynthetic electron transport chain. Direct measurement of the turnover time of photosynthetic hydrogen production (Greenbaum, 1979 and 1982) demonstrated that this parameter was comparable to the turnover time of oxygen production. It was also shown (Greenbaum, 1988) that net conversion efficiencies of 5-10% could be achieved in the linear low-intensity region of the light saturation curve.

## Research Problems

The hydrogenase enzyme is synthesized *de novo* under anaerobic conditions. In normal

photosynthesis carbon dioxide is the preferred electron acceptor for photogenerated reductant from Photosystem I. However, direct kinetic competition between hydrogen evolution and the Calvin cycle can easily be observed (Graves et al. 1989; Cinco et al. 1993). The three scientific research problems associated with photosynthetic hydrogen and oxygen production are (1) oxygen sensitivity of hydrogenase; (2) antenna size, bicarbonate and the light saturation problem; and (3) the minimum number of light reactions required to split water to molecular hydrogen and oxygen.

### ***Oxygen Sensitivity of Hydrogenase***

In the application of intact unicellular green algae for hydrogen production one is confronted with the problem of oxygen sensitivity of the hydrogenase enzyme. Hydrogenase is synthesized under anaerobic conditions and, at present, must be kept that way in order to preserve its functionality. In one approach, oxygen and hydrogen by green algae are coproduced in the same volume. Therefore, a way must be found to prevent inhibition of hydrogenase activity by the photosynthetically produced oxygen. This challenging problem is the focus of research at the National Renewable Energy Laboratory (Ghirardi et al., 1997).

### ***Antenna Size, Bicarbonate and the Light Saturation Problem***

In full sunlight,  $\approx 1000 \text{ W/m}^2$ , there exists a kinetic imbalance between the rate of photon excitation of the reaction centers and the ability of the thermally-activated electron transport chains to process photogenerated electrons. Whereas the reaction centers can receive photoexcitations at the rate of  $\approx 2000 \text{ sec}^{-1}$ , movement through the electron transport chain is of the order of  $200 \text{ sec}^{-1}$  or less (Gibbs et al., 1973). Therefore, normal photosynthesis saturates at much less than full sunlight, typically  $\approx 10\%$ .

Since there is little opportunity to increase the rate of thermally-activated electrons through the photosynthetic electron transport chain, an alternate strategy is to reduce the antenna size. Kinetic balance between the rate of photon excitation and rate of photosynthetic electrons can, in principle, be balanced, even at full sunlight, by reducing the absolute antenna size per reaction center. If such a response could be achieved in a real-world system, photosynthetic productivity on a per chlorophyll basis would increase and high solar irradiances would be converted to useful biomass energy. Linearization of the light saturation curve of photosynthesis was demonstrated by Herron and Mauzerall (1972). Melis et al. (1998) have demonstrated linearization of the light saturation curve for high-light grown cultures of *Dunaliella*. These results indicate that the concept is technically correct.

An additional complication of the light saturation problem involves the requirement of bicarbonate to optimize electron transport through Photosystem II. Since carbon dioxide/bicarbonate are the exclusive sink for photosynthetically generated reductant they need to be removed so that the flow of electrons produces hydrogen rather than carbon dioxide fixation compounds. Complete removal of carbon dioxide, however, impairs electron transport in Photosystem II and further reduces the saturating light intensity for sustained simultaneous photoproduction of hydrogen and oxygen by about factor of 10. Qualitatively speaking, light saturation occurs at about  $10 \text{ W/m}^2$ . One strategy to overcome this limitation is to take advantage of the differential affinity of  $\text{CO}_2/\text{bicarbonate}$  between the Photosystem II binding site and the Calvin cycle. Such an approach has been explored by Cinco et al. (1993) in which light-activated hydrogen and oxygen evolution as a function of  $\text{CO}_2$  concentration in helium were measured for the unicellular green alga *Chlamydomonas reinhardtii*. The concentrations were 58, 30, 0.8 and 0 ppm  $\text{CO}_2$ . The objective of these experiments was to study the differential affinity of  $\text{CO}_2/\text{HCO}_3^-$  for their respective Photosystem II and Calvin cycle binding sites vis-à-vis photoevolution of molecular oxygen and the competitive pathways of hydrogen photoevolution and  $\text{CO}_2$  photoassimilation. The maximum rate of hydrogen evolution occurred at 0.8 ppm  $\text{CO}_2$ . The key result of this work was that the rate of photosynthetic hydrogen evolution can be increased, at least partially, by satisfying the Photosystem II  $\text{CO}_2/\text{HCO}_3^-$  binding site requirement without fully activating the Calvin-Benson  $\text{CO}_2$  reduction pathway. These preliminary experiments suggest that mutants of *Chlamydomonas reinhardtii* that have a genetically engineered low  $\text{CO}_2$  affinity for the

Calvin cycle and relatively higher affinity for the PS II  $\text{CO}_2/\text{HCO}_3^-$  binding site may be good candidates to explore for relieving the  $\text{CO}_2$  part of the light saturation constraint.

### **Thermodynamic Driving Pressure of Photosynthetic Hydrogen Production**

We have shown that the thermodynamic driving pressure of hydrogen production in the green alga *Scenedesmus* D<sub>3</sub> is equal to or greater than one atmosphere. This was accomplished by measuring the rate of photosynthetic oxygen production by *Scenedesmus* in one-atmosphere of pure hydrogen. The practical significance of this work is that it helps to minimize the amount of pump work required to deliver hydrogen at a useable pressure. At a minimum, the amount of energy saved is  $\Delta W = RT \ln(P_{\text{final}}/P_{\text{initial}})$ . Since this is the reversible equilibrium thermodynamic value, the actual value for real gases, including irreversible processes, will be several times this. The amount of energy saved can be calculated to be in the range 68,500 - 114,200 J mol<sup>-1</sup>.  $\Delta G_p^\circ$  for the reaction,  $\text{H}_2\text{O} \rightarrow \text{H}_2 + \frac{1}{2}\text{O}_2$ , is 237,200 J mol<sup>-1</sup>. Another significant aspect of this result is that it demonstrates for the first time that the oxygen evolution enzyme is insensitive to the presence of high concentrations (*i.e.*, pure) hydrogen. That is to say, unlike the well-known sensitivity of the hydrogenase enzyme to even low concentrations of oxygen, the oxygen evolving complex of photosynthesis is unaffected by the presence of hydrogen. The indirect method of measuring oxygen was necessary because it is experimentally impossible to measure photosynthetically produced hydrogen against a background carrier gas of pure hydrogen since the gas sensitive semiconducting detectors that are used for the hydrogen analysis are saturated in pure hydrogen. Isotopic labeling experiments are currently under way for direct measurement of hydrogen production.

[Note: Much of the text of this abstract has been previously published. Please see E. Greenbaum and J. W. Lee, "Photosynthetic Hydrogen and Oxygen Production by Green Algae: An Overview," in *BioHydrogen*, O. Zaborsky, Ed. pp.235-242 (1998). [Copyright Plenum Press, New York, 1998]

### **References**

- Bishop NI and Gaffron H (1963) On the interrelation of the mechanisms for oxygen and hydrogen evolution in adapted algae. In *Photosynthetic Mechanisms in Green Plants*, B. Kok and A. T. Jagendorf, Eds. (Natl. Acad. Sci.-Natl. Res. Council, Washington, DC), pp. 441-451
- Bishop NI, Fricke M, and Jones LW (1977) Photohydrogen production in green algae: water serves as the primary substrate for hydrogen and oxygen production. In *Biological Solar Energy Production*, A. Mitsui, S. Miyachi, A. San Pietro, and S. Tamura, Eds. (Academic Press, New York) pp. 1-22
- Cinco RM, MacInnis JM and Greenbaum E (1993) The Role of Carbon Dioxide in Light-Activated Hydrogen Production by *Chlamydomonas reinhardtii*. *Photosyn. Res.* 38: 27-33 (1993)
- Gaffron H (1960) Energy storage: photosynthesis. In *Plant Physiology* **1B**: 4-277
- Gaffron H and Rubin J (1942) Fermentative and photochemical production of hydrogen in algae. *J. Gen. Physiol.* 26: 219-240
- Ghirardi ML, Togasaki RK and Seibert M (1997) Oxygen sensitivity of algal hydrogen production. *Appl. Biochem. Biotech.*, in press
- Gibbs M, Hollaender A, Kok B, Krampitz, LO and San Pietro A (1973) Proceedings of the Workshop on Bio-Solar Conversion, September 5-6, 1973, Bethesda, MD, NSF RANN Grant GI 40253 to Indiana University
- Graves DA, Tevault CV and Greenbaum E (1989) Control of Photosynthetic Reductant: The Effect of Light and Temperature on Sustained Hydrogen Photoevolution in *Chlamydomonas* sp. in an Anoxic Carbon Dioxide Containing Atmosphere. *Photochem. Photobiol.* 50: 571-576
- Greenbaum E (1982) Photosynthetic hydrogen and oxygen production: kinetic studies. *Science* 215: 291-293
- Greenbaum E (1977a) The photosynthetic unit size of hydrogen evolution. *Science* 196: 878-879
- Greenbaum E (1977b) The molecular mechanisms of photosynthetic hydrogen and oxygen production. In *Biological Solar Energy Production*, A. Mitsui, S. Miyachi, A. San Pietro, and S. Tamura, Eds. (Academic Press, New York) pp. 101-107
- Greenbaum E (1979) The turnover times and pool sizes of photosynthetic hydrogen production by green algae. *Solar Energy* 23: 315-320
- Greenbaum E (1988) Energetic efficiency of hydrogen photoevolution by algal water splitting. *Biophys. J.* 54: 365-368

- Greenbaum E (1980) Simultaneous photoproduction of hydrogen and oxygen by photosynthesis. *Biotechnol. Bioengin. Symp.* 10, 1-13
- Greenbaum E, Lee JW, Blankinship SL and Tevault CV (1997b) Hydrogen and oxygen production in mutant Fud26 of *Chlamydomonas reinhardtii* Proceedings of the 1997 U.S. DOE Hydrogen Program Review. May 21-23, 1997, Herndon, VA, in press.
- Greenbaum E, Lee JW, Tevault CV, Blankinship SL, Mets LJ, and Owens TG (1997a) Photosystem I measurement in mutants B4 and F8 of *C. reinhardtii*. *Science* 277: 167-168, Nature, in press.
- Greenbaum E, Lee JW, Tevault CV, Blankinship SL and Mets LJ (1995) CO<sub>2</sub> Fixation and photoevolution of H<sub>2</sub> and O<sub>2</sub> in a mutant of *Chlamydomonas* lacking Photosystem I. *Nature* 376: 438-441
- Herron HA and Mauzerall D. (1972) The development of photosynthesis in a greening mutant of *Chlorella* and an analysis of the light saturation curve. *Plant Physiol.* 50: 141-148
- Kaltwasser H, Stuart TS and Gaffron H (1969) Light-Dependent Hydrogen Evolution by *Scenedesmus*. *Planta* (Berlin) 89: 309-322.
- Lee JW, Tevault CV, Owens TG and Greenbaum E (1996) Oxygenic photoautotrophic growth without Photosystem I. *Science* 273: 364-367
- Lien S and San Pietro A (1975) An Inquiry into Biophotolysis of Water to Produce Hydrogen, NSF RANN report under grant GI 40253 to Indiana University
- Melis A, Neidhardt J, Baroli I, and Benemann, JR (1998) Maximizing photosynthetic productivity and light utilization in microalgae by minimizing the light-harvesting chlorophyll antenna size of the photosystems. These proceedings, in press
- Mitsui A, Miyachi S, San Pietro A and Tamura S, Eds. (1977) *Biological Solar Energy Production* (Academic Press, New York)
- Spruit CJP (1958) Simultaneous photoproduction of hydrogen and oxygen by *Chlorella*. *Mededel. Landbouwhogeschool* (Wageningen/Nederland) 58:1-17
- Stuart TS and Gaffron H (1972) The mechanism of hydrogen photoproduction by several algae. *Planta* (Berlin) 106: 91-100
- Stuart TS and Gaffron H (1971) The kinetics of hydrogen photoproduction by adapted *Scenedesmus*. *Planta* (Berlin) 100: 228-243